

TrkA+B+C Rabbit mAb

货号: **AYM30058**

产品信息

| | |
|-------|---|
| 反应 | Human,Mouse,Rat |
| 宿主 | Rabbit |
| 克隆性 | Monoclonal |
| 预测反应 | |
| 应用 | WB IHC ICC |
| 推荐浓度 | WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 ICC: 1:50 - 1:200 |
| 理论分子量 | 140kDa |
| 实测分子量 | 140kDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | SH-SY5Y,Mouse brain,Rat brain |
| 细胞定位 | axon,dendrite,early endosome,late endosome,plasma membrane |
| 纯化 | Affinity purification |

抗原信息


| | |
|------|---|
| 抗原信息 | Recombinant fusion protein corresponding to Human TrkA+B+C. |
|------|---|

靶点信息


| | |
|-------|----------------------|
| 研究背景 | |
| 基因ID | 4914,4916,4915 |
| 基因名 | NTRK1,NTRK3,NTRK2 |
| Swiss | P04629/Q16288/Q16620 |

| | |
|----|----------|
| 别名 | TrkA+B+C |
|----|----------|

产品验证



Western blot analysis of TrkA+B+C expressed in SH-SY5Y, Mouse brain, Rat brain using TrkA+B+C Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.



Immunohistochemical analysis of paraffin-embedded mouse brain, using TrkA+B+C Antibody.

实验步骤

访问官网浏览详情: www.ablybio.cn