

— ABLYBIO, Help Your Research



Somatostatin Receptor 2 Rabbit mAb

货号: **AYM30088**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IP
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IP: 1:20 - 1:50
理论分子量	40kDa/41kDa
实测分子量	41kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	MCF7,C6,SH-SY5Y,Neuro-2a,MDA-MB-231,Mouse thymus
细胞定位	Cell membrane,Cytoplasm,Multi-pass membrane protein
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human Somatostatin Receptor 2.
------	--

靶点信息

研究背景	Somatostatin acts at many sites to inhibit the release of many hormones and other secretory proteins. The biologic effects of somatostatin are probably mediated by a family of G protein-coupled receptors that are expressed in a tissue-specific manner. SSTR2 is a member of the superfamily of receptors having seven transmembrane segments and is expressed in highest levels in cerebrum and kidney.
基因ID	6752
基因名	SSTR2
Swiss	P30874
别名	Somatostatin Receptor 2,Somatostatin Receptor 2 Rabbit mAb,SSTR2,SRIF-1

产品验证



Western blot analysis of Somatostatin Receptor 2 expressed in Mouse thymus, MCF7, C6 using Somatostatin Receptor 2 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/>www.ablybio.cn)