

ACVR1B Rabbit mAb

货号: **AYM30107**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	52kDa
实测分子量	52kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,HepG2,U-87MG,Mouse brain,Rat brain
细胞定位	cell surface,cytosol,plasma membrane
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human ACVR1B.
序列	MAESAGASSFFPLVLLLAGSGGSGPRGVQALLCACTSLQANYTCETDGACMVSIFNLDGMEHHVRTCIPKVELVPAGK PFYCLSEDLRNTHCCYTDY

靶点信息

研究背景	This gene encodes an activin A type IB receptor. Activins are dimeric growth and differentiation factors which belong to the transforming growth factor-beta (TGF-beta) superfamily of structurally related signaling proteins. Activins signal through a heteromeric complex of receptor serine kinases which include at least two type I and two type II receptors. This protein is a type I receptor which is essential for signaling. Mutations in this gene are associated with pituitary tumors. Alternate splicing results in multiple transcript variants.[provided by RefSeq, Jun 2010]
------	--

基因ID	91
基因名	ACVR1B
Swiss	P36896
别名	ACTRIB; ACVRLK4; ALK4; SKR2

产品验证



Western blot analysis of ACVR1B expressed in Mouse brain, Rat brain, 293T using ACVR1B Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn