

# BAG2 Rabbit mAb

货号: **AYM30111**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IF/ICC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IF/ICC:</b> 1:50 - 1:200
理论分子量	24kDa
实测分子量	24kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	A-549,PC-3,Jurkat
细胞定位	axon,cytosol,dendrite,dendritic microtubule
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein corresponding to Human BAG2.
序列	MAQAKINAKANEGRFRCSSSMADRSSRLLESLDQLELRVEALREAATAVEQEKEILLEMHSIQNSQDMRQISDGEREELNL TANRLMGRTLTVESVET

## 靶点信息

研究背景	BAG proteins compete with Hip for binding to the Hsc70/Hsp70 ATPase domain and promote substrate release. All the BAG proteins have an approximately 45-amino acid BAG domain near the C terminus but differ markedly in their N-terminal regions. The predicted BAG2 protein contains 211 amino acids. The BAG domains of BAG1, BAG2, and BAG3 interact specifically with the Hsc70 ATPase domain in vitro and in mammalian cells. All 3 proteins bind with high affinity to the ATPase domain of Hsc70 and inhibit its chaperone activity in a Hip-repressible manner. [provided by RefSeq, Jul 2008]
------	---

基因ID	9532
基因名	BAG2
Swiss	O95816
别名	BAG-2; dj41711.2

## 产品验证



Western blot analysis of BAG2 expressed in A-549,PC-3,Jurkat using BAG2 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)