

# CD89 Rabbit mAb

货号: **AYM30119**

## 产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC IP FC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IHC:</b> 1:50 - 1:200 <b>IF/ICC:</b> 1:50 - 1:200 <b>IP:</b> 1:20 - 1:50 <b>FC:</b> 1:20 - 1:50
理论分子量	19-32kDa
实测分子量	70kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T
细胞定位	Cell membrane,Secreted,Single-pass type I membrane protein
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein corresponding to Human CD89.
序列	LMIKNSTYREIGRRLKFWNETDPEFVIDHMDANKAGRYQCQYRIGHYFRYSDTLELVVTGLYGKPFLSADRGLVLMPEGNISLTCSSAHIPFDRFSLAKEGELSLPQHQSGEHPANFSLGPVDLNVSGIYRCYGWYNRSPYLWSFSPNALELVVTDSDSIHQDYTTQNLIRMAVAGLVLLVALLAILVENWHSHTALNKEASADVAEPSWSQMCQPGLTFARTPSVCK

## 靶点信息

研究背景	This gene is a member of the immunoglobulin gene superfamily and encodes a receptor for the Fc region of IgA. The receptor is a transmembrane glycoprotein present on the surface of myeloid lineage cells such as neutrophils, monocytes, macrophages, and eosinophils, where it mediates immunologic responses to pathogens. It interacts with IgA-opsonized targets and triggers several immunologic defense processes, including phagocytosis, antibody-dependent cell-mediated cytotoxicity, and stimulation of the release of inflammatory mediators. Multiple alternatively spliced transcript variants encoding different isoforms have been described for this gene.
基因ID	2204
基因名	FCAR
Swiss	P24071
别名	FCAR;CD89;CTB-61M7.2;FcalphaRI

## 产品验证



Western blot analysis of CD89 expressed in 293T using CD89 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)