

CD80 Rabbit mAb

货号: **AYM30122**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200
理论分子量	18kDa/29kDa/33kDa
实测分子量	60kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Rat lung
细胞定位	Membrane,Single-pass type I membrane protein
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human CD80.
序列	MGHTRRQGTSPSKCPYLNFFQLLVLAGLSHFCSGVIVHTKEVKEVATLSCGHNVSVLELAQTRIWQKEKKMLTMMMSG DMNIWPEYKNRTIFDITNNLS

靶点信息

研究背景	The protein encoded by this gene is a membrane receptor that is activated by the binding of CD28 or CTL A-4. The activated protein induces T-cell proliferation and cytokine production. This protein can act as a receptor for adenovirus subgroup B and may play a role in lupus neuropathy.
基因ID	941

基因名	CD80
Swiss	P33681
别名	B7;B7-1;B7.1;BB1;CD28LG;CD28LG1;LAB7;CD80

产品验证



Western blot analysis of CD80 expressed in Rat lung using CD80 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn