

Intra Acrosomal Protein Rabbit mAb

货号: **AYM30124**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	9kDa/13-28kDa
实测分子量	14-36kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Rat testis
细胞定位	Cytoplasmic vesicle,acrosome,secretory vesicle
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human Intra Acrosomal Protein.
序列	QPNELSGSIDHQTSVQQLPGEFFSLENPSDAEALYETSSGLNTLSEHGSSEHGSSKHTVAEHTSGEHAESEHASGEPAAEHAEGEHTVGEQPSGEGPSGEHLSGEQPLSELESGEQPSDEQPSGEGHSGEQPSGEGQASGEGPSGEHASGEQASGAPISSTSTGTILNCYTCAYMNDQKCLRGEGETCITQNSQQCMLKKIFEGGKLQFMVQGCENMCPMSMNLFSHGTRMQIICCRNQSFCNKI

靶点信息

研究背景	This gene encodes a testis-specific, differentiation antigen, acrosomal vesicle protein 1, that arises within the acrosomal vesicle during spermatogenesis, and is associated with the acrosomal membranes and matrix of mature sperm. The acrosomal vesicle protein 1 may be involved in sperm-zona binding or penetration. Alternatively spliced transcript variants have been described.
基因ID	56

基因名	ACRV1
Swiss	P26436
别名	ACRV1;D11S4365;SP-10;SPACA2

产品验证



Western blot analysis of Intra Acrosomal Protein expressed in Rat testis using Intra Acrosomal Protein Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/protein s: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn