

# PFDN5 Rabbit mAb

货号: **AYM30125**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IP
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IHC:</b> 1:50 - 1:200 <b>IP:</b> 1:20 - 1:50
理论分子量	7kDa/12kDa/17kDa
实测分子量	17kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Raji,U-87MG,Mouse brain,Rat brain
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein corresponding to Human PFDN5.
序列	MAQSINITELNLPQLEMLKNQLDQEVFLSTLSIAQLKVVQTKYVEAKDCLNVLNKSNEGKELLVPLTSSMYVPGKLHDVEHV LIDVGTGYVVEKTAEDAKDFFKRKIDFLTKQMEKIQPALQEKHAMKQAVMEMMSQKIQQLTALGAAQATAKA

## 靶点信息

研究背景	This gene encodes a member of the prefoldin alpha subunit family. The encoded protein is one of six sub units of prefoldin, a molecular chaperone complex that binds and stabilizes newly synthesized polypeptid es, thereby allowing them to fold correctly. The complex, consisting of two alpha and four beta subunits, forms a double beta barrel assembly with six protruding coiled-coils. The encoded protein may also repre ss the transcriptional activity of the proto-oncogene c-Myc. Alternatively spliced transcript variants encodi ng different isoforms have been described.
------	--

基因ID	5204
基因名	PFDN5
Swiss	Q99471
别名	PFDN5;MM-1;MM1;PFD5

## 产品验证



Western blot analysis of PFDN5 expressed in Mouse brain,Rat brain,Raji using PFDN5 Rabbit mAb at 1:1 000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)