

IL11RA Rabbit mAb

货号: **AYM30128**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC FC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 FC: 1:20 - 1:50
理论分子量	41kDa/45kDa
实测分子量	45kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse kidney,Mouse heart,Mouse skeletal muscle,Rat liver
细胞定位	Membrane,Single-pass type I membrane protein
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human IL11RA.
序列	WGPPGVQYQPGRSVKLCCPGVTAGDPVSWFRDGEKLLQGPDSGLGHELVLQAQADSTDEGTYICQTLDGALGGTVTL QLGYPPARPVVSCQAADYENFCTWSPSQISGLPTRYLTSYRKKTVLGADSQRRSPSTGPWPCPDPLGAARCVVHGAE FWSQYRINVTEVNPGLGASTRLLDVSLQILRPDPPQGLRVESVPGYPRRLRASWTPASWPCQPHFLKFRLLQYRPAQHPA WSTVEPAGLEEVITDAVAGLPHA

靶点信息

研究背景	Interleukin 11 is a stromal cell-derived cytokine that belongs to a family of pleiotropic and redundant cytokines that use the gp130 transducing subunit in their high affinity receptors. This gene encodes the IL-11 receptor, which is a member of the hematopoietic cytokine receptor family. This particular receptor is very similar to ciliary neurotrophic factor, since both contain an extracellular region with a 2-domain structure composed of an immunoglobulin-like domain and a cytokine receptor-like domain. Multiple alternatively spliced transcript variants have been found for this gene.
基因ID	3590
基因名	IL11RA
Swiss	Q14626
别名	IL11RA;CRSDA

产品验证



Western blot analysis of IL11RA expressed in Mouse kidney, Mouse heart, Mouse skeletal muscle using IL11RA Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn