— ABLYBIO, Help Your Research

VEGFD Rabbit mAb

货号: AYM30143

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC FC
推荐浓度	**WB:** 1:500 - 1:2000 **IHC:** 1:50 - 1:200 **IF/ICC:** 1:50 - 1:200 **FC:** 1:20 - 1:50
理论分子量	40kDa
实测分子量	40kDa
形式	Liquid
保存条件	Store at -20℃. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	NCI-H460
细胞定位	Secreted
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human VEGFD.

靶点信息

研究背景	The protein encoded by this gene is a member of the platelet-derived growth factor/vascular endothelial growth factor (PDGF/VEGF) family and is active in angiogenesis, lymphangiogenesis, and endothelial cell growth. This secreted protein undergoes a complex proteolytic maturation, generating multiple processe d forms which bind and activate VEGFR-2 and VEGFR-3 receptors. This protein is structurally and function ally similar to vascular endothelial growth factor C. Read-through transcription has been observed betwe en this locus and the upstream PIR (GeneID 8544) locus.
基因ID	2277
基因名	VEGFD
Swiss	O43915 (https://www.uniprot.org/uniprotkb/O43915/entry)
别名	VEGFD,VEGFD Rabbit mAb,c-Fos-induced growth factor,FIGF

产品验证



Western blot analysis of VEGFD expressed in NCI-H460 using VEGFD Rabbit mAb at 1:1000. Secondary an tibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% no n-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (https://www.ablybio.cn//www.ablybio.cn)