

R Cadherin Rabbit mAb

货号: **AYM30147**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	92kDa/100kDa
实测分子量	130kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HepG2,Mouse brain,Mouse thymus,Rat brain,Rat skeletal muscle
细胞定位	Cell membrane,Single-pass type I membrane protein
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human R Cadherin.
序列	KRREKERHTKQLLIDPEDDVRDNILKYDEEGGGEDQDYDLSQLQQPEAMGHVPSKAPGVERRVDERPVGAEPQYPIRPM VPHPGDIGDFINEGLRAADNDPTAPPYDSSLVFDYEGSGSTAGSVSSLNSSSSGDQDYDYLNDWGPRFKKLADMYGGG EED

靶点信息

研究背景	This gene is a classical cadherin from the cadherin superfamily. The encoded protein is a calcium-dependent cell-cell adhesion glycoprotein comprised of five extracellular cadherin repeats, a transmembrane region and a highly conserved cytoplasmic tail. Based on studies in chicken and mouse, this cadherin is thought to play an important role during brain segmentation and neuronal outgrowth. In addition, a role in kidney and muscle development is indicated. Of particular interest are studies showing stable cis-heterodimers of cadherins 2 and 4 in cotransfected cell lines. Previously thought to interact in an exclusively homophilic manner, this is the first evidence of cadherin heterodimerization. Three transcript variants encoding different isoforms have been found for this gene.
基因ID	1002
基因名	CDH4
Swiss	P55283
别名	CDH4;CAD4;R-CAD;RCAD

产品验证



Western blot analysis of R Cadherin expressed in Mouse brain, Rat brain, HepG2 using R Cadherin Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn