

BACH1/BRIP1 Rabbit mAb

货号: **AYM30154**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	112kDa/140kDa
实测分子量	105kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Jurkat,MCF7
细胞定位	Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human BACH1/BRIP1.
序列	MSSMWSEYTIGGVKIYFPYKAYPSQLAMMNSILRGLNSKQHCLLESPTGSGKSLALLCSALAWQQSLSGKPADEGVSEKA EVQLSCCCACHSKDFTNNDMNQGTSRHFNYSTPPSERNGTSSTCQDSPEKTTAAKLSAKKQASIYRDENDDFQVEKK RIRPLETTQIRKRHCFGTEVHNLDAKVDGKTVKLNPLEKINSFSPQKPPGHCSRCCCSTKQGNSQESSNTIKKDHTGK SKIPKIYFGTRTHKQIAQIT

靶点信息

研究背景	The protein encoded by this gene is a member of the RecQ DEAH helicase family and interacts with the B RCT repeats of breast cancer, type 1 (BRCA1). The bound complex is important in the normal double-strand break repair function of breast cancer, type 1 (BRCA1). This gene may be a target of germline cancer-inducing mutations.
基因ID	83990

基因名	BRIP1
Swiss	Q9BX63
别名	BRIP1;BACH1;FANCJ;OF

产品验证



Western blot analysis of BACH1/BRIP1 expressed in Jurkat, MCF7 using BACH1/BRIP1 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn