

Hsp22 Rabbit mAb

货号: **AYM30155**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	21kDa
实测分子量	22kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	RD,Mouse liver,Mouse heart,Rat heart
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human Hsp22.
序列	MADGQMPFSCHYPSRLRRDPFRD SPLSRLDDGFGMDPFPDDL TASWPDWALPRLSSAWPGTLRSGMVPRGPTATA RFGVPAEGRTPPFPGEWPKVCVNVHSFKPEELMVKTKDGYVEVSGKHEEKQEGGIVSKNFTKKIQLPAEVD PVTVFAS LSPEGLLIIEAPQVPPYSTFGESSFNNELPQDSQEVTCT

靶点信息

研究背景	The protein encoded by this gene belongs to the superfamily of small heat-shock proteins containing a conserved alpha-crystallin domain at the C-terminal part of the molecule. The expression of this gene is induced by estrogen in estrogen receptor-positive breast cancer cells, and this protein also functions as a chaperone in association with Bag3, a stimulator of macroautophagy. Thus, this gene appears to be involved in regulation of cell proliferation, apoptosis, and carcinogenesis, and mutations in this gene have been associated with different neuromuscular diseases, including Charcot-Marie-Tooth disease.
------	---

基因ID	26353
基因名	HSPB8
Swiss	Q9UJY1
别名	HSPB8;CMT2L;DHMN2;E2IG1;H11;HMN2;HMN2A;HSP22

产品验证



Western blot analysis of Hsp22 expressed in Mouse liver,Rat heart,RD using Hsp22 Rabbit mAb at 1:10 00. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Bl ocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn