

Smac / Diablo Rabbit mAb

货号: **AYM30213**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC IP FC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 IP: 1:20 - 1:50 FC: 1:20 - 1:50
理论分子量	21kDa/22kDa/27kDa
实测分子量	21kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	SW620,BT-474,Mouse spleen
细胞定位	Mitochondrion
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human Smac / Diablo.
序列	AVPIAQKSEPHLSSEALMRRRAVSLVTDSTSTFLSQTTYALIEAITEYTKAVYTLTSLYRQYTSLLGKMNSEEEDEVWQVIIGA RAEMTSKHQEYLKLETTWMTAVGLSEMAAEAAAYQTGADQASITARNHIQLVKLQVEEVHQLSRKAETKLAEAQIEELRQKT QEEGEERAESEQEAYLRED


靶点信息

研究背景	This gene encodes an inhibitor of apoptosis protein (IAP)-binding protein. The encoded mitochondrial protein enters the cytosol when cells undergo apoptosis, and allows activation of caspases by binding to inhibitor of apoptosis proteins. Overexpression of the encoded protein sensitizes tumor cells to apoptosis. A mutation in this gene is associated with young-adult onset of nonsyndromic deafness-64. Alternative splicing results in multiple transcript variants encoding different isoforms.
基因ID	56616
基因名	DIABLO
Swiss	Q9NR28
别名	DIABLO;DFNA64;SMAC

产品验证



Immunohistochemical analysis of paraffin-embedded human kidney, using Smac/Diablo Antibody.



Western blot analysis of Smac / Diablo expressed in SW620,BT-474,Mouse spleen using Smac / Diablo Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn