— ABLYBIO, Help Your Research

PAK1 Rabbit mAb

货号: AYM30219

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IP FC ICC/IF
推荐浓度	**WB:** 1:500 - 1:2000 **IHC:** 1:50 - 1:200 **IF/ICC:** 1:50 - 1:200 **IP:** 1:20 - 1:50 **FC:** 1:20 - 1:50
理论分子量	60kDa/61kDa
实测分子量	66kDa
形式	Liquid
保存条件	Store at -20℃. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	K-562,C6
细胞定位	Cell junction,Cell membrane,Cell projection,Cytoplasm,focal adhesion,invadopodium,ruffle membrane
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human PAK1.


靶点信息

研究背景	This gene encodes a family member of serine/threonine p21-activating kinases, known as PAK proteins. These proteins are critical effectors that link RhoGTPases to cytoskeleton reorganization and nuclear signaling, and they serve as targets for the small GTP binding proteins Cdc42 and Rac. This specific family member regulates cell motility and morphology. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.
基因ID	5058
基因名	PAK1
Swiss	Q13153 (https://www.uniprot.org/uniprotkb/Q13153/entry)
别名	PAK1,PAK1 Rabbit mAb,Alpha-PAK,p21-activated kinase 1,p65-PAK

产品验证



Western blot analysis of PAK1 expressed in K-562,C6 using PAK1 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.



Immunofluorescence analysis of U2OS cells labelling PAK1 with . The cells were fixed with cold 100% methanol (10min, 4℃) and blocked in 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween 20 for 1h. The cells were then incubate with Goat Anti-Rabbit IgG Fluor488 (1:200，shown in green) at room temprature for 1h.

实验步骤

访问官网浏览详情: www.ablybio.cn (https://www.ablybio.cn//www.ablybio.cn)