

STC1 Rabbit mAb

货号: **AYM30236**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IP
推荐浓度	WB: 1:500 - 1:2000 IP: 1:20 - 1:50
理论分子量	20kDa/27kDa
实测分子量	31-37kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse lung
细胞定位	Secreted
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human STC1.
序列	THEAEQNDSVSPRKSrvAAQNSAEVVRCLNSALQVGCgAFACLENSTCDTDGMYDICKSFLYSAAKFDTQGKAFVKESL KCIANGVTSKVF LAIRRCSTFQRMIAEVQEECYSKLNVCSIAKRNPEAITEVVQLPNHFSNRYNRLVRSLLCEDTVSTIRD SLMEKIGPNMASLFHILQTDHCAQTHPRADFNRRRTNEPQKLKVLRLNLRGEEDSPSHIKRTSHESA

靶点信息

研究背景	This gene encodes a secreted, homodimeric glycoprotein that is expressed in a wide variety of tissues and may have autocrine or paracrine functions. The gene contains a 5' UTR rich in CAG trinucleotide repeats. The encoded protein contains 11 conserved cysteine residues and is phosphorylated by protein kinase C exclusively on its serine residues. The protein may play a role in the regulation of renal and intestinal calcium and phosphate transport, cell metabolism, or cellular calcium/phosphate homeostasis. Overexpression of human stanniocalcin 1 in mice produces high serum phosphate levels, dwarfism, and increased metabolic rate. This gene has altered expression in hepatocellular, ovarian, and breast cancers.
基因ID	6781
基因名	STC1
Swiss	P52823
别名	STC1;STC

产品验证



Western blot analysis of STC1 expressed in Mouse lung using STC1 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn