

— ABLYBIO, Help Your Research



WASL Rabbit mAb

货号: **AYM30243**

产品信息

| | |
|-------|--|
| 反应 | Human,Mouse |
| 宿主 | Rabbit |
| 克隆性 | Monoclonal |
| 预测反应 | |
| 应用 | WB IHC IF/ICC FC |
| 推荐浓度 | WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 FC: 1:20 - 1:50 |
| 理论分子量 | 54kDa |
| 实测分子量 | 55kDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | HeLa,U-87MG,A-549,Mouse brain,Mouse lung,Rat brain |
| 细胞定位 | Cytoplasm,Nucleus,cytoskeleton |
| 纯化 | Affinity purification |


抗原信息

| | |
|------|---|
| 抗原信息 | Recombinant fusion protein corresponding to Human WASL. |
|------|---|

靶点信息

| | |
|-------|--|
| 研究背景 | This gene encodes a member of the Wiskott-Aldrich syndrome (WAS) protein family. Wiskott-Aldrich syndrome proteins share similar domain structure, and associate with a variety of signaling molecules to alter the actin cytoskeleton. The encoded protein is highly expressed in neural tissues, and interacts with several proteins involved in cytoskeletal organization, including cell division control protein 42 (CDC42) and the actin-related protein-2/3 (ARP2/3) complex. The encoded protein may be involved in the formation of long actin microspikes, and in neurite extension. |
| 基因ID | 8976 |
| 基因名 | WASL |
| Swiss | O00401 |
| 别名 | WASL,WASL Rabbit mAb,Neural Wiskott-Aldrich syndrome protein |

产品验证



Western blot analysis of WASL expressed in Mouse brain, HeLa, U-87MG using WASL Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/>www.ablybio.cn)