— ABLYBIO, Help Your Research

MYO7A Rabbit mAb

货号: AYM30265

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IF/ICC FC
推荐浓度	**WB:** 1:500 - 1:2000 **IF/ICC:** 1:50 - 1:200 **FC:** 1:20 - 1:50
理论分子量	138kDa/240kDa/249kDa/250kDa/254kDa
实测分子量	254kDa
形式	Liquid
保存条件	Store at -20℃. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse testis,Rat testis
细胞定位	Cytoplasm,cell cortex,cytoskeleton
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human MYO7A.

靶点信息

研究背景	This gene is a member of the myosin gene family. Myosins are mechanochemical proteins characterized by the presence of a motor domain, an actin-binding domain, a neck domain that interacts with other pro teins, and a tail domain that serves as an anchor. This gene encodes an unconventional myosin with a ve ry short tail. Defects in this gene are associated with the mouse shaker-1 phenotype and the human Ushe r syndrome 1B which are characterized by deafness, reduced vestibular function, and (in human) retinal degeneration. Alternative splicing results in multiple transcript variants.
基因ID	4647
基因名	MYO7A
Swiss	Q13402 (https://www.uniprot.org/uniprotkb/Q13402/entry)
别名	MYO7A,MYO7A Rabbit mAb,USH1B

产品验证



Western blot analysis of MYO7A expressed in Mouse testis,Rat testis using MYO7A Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (https://www.ablybio.cn//www.ablybio.cn)