

PIAS1+PIAS2 Rabbit mAb

货号: **AYM30276**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC IP FC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 IP: 1:20 - 1:50 FC: 1:20 - 1:50
理论分子量	76kDa
实测分子量	76kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HepG2,Mouse brain
细胞定位	nuclear speck,nucleoplasm,nucleus,PML body
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human PIAS1+PIAS2.
------	--

靶点信息


研究背景	This gene encodes a member of the protein inhibitor of activated STAT (PIAS) family. PIAS proteins function as SUMO E3 ligases and play important roles in many cellular processes by mediating the sumoylation of target proteins. This protein plays a central role as a transcriptional coregulator of numerous cellular pathways including the STAT1 and nuclear factor kappaB pathways. Alternate splicing results in multiple transcript variants. [provided by RefSeq, Mar 2016]
基因ID	8554, 9063

基因名	PIAS1
Swiss	O75925/O75928
别名	PIAS1+PIAS2

产品验证



Western blot analysis of PIAS1+PIAS2 expressed in HepG2, Mouse brain using PIAS1+PIAS2 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.



Immunohistochemical analysis of paraffin-embedded human colon, using PIAS1+PIAS2+PIAS3 Antibody.

实验步骤

访问官网浏览详情: www.ablybio.cn