

EHMT2/G9A Rabbit mAb

货号: **AYM30277**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200
理论分子量	160,170kDa
实测分子量	170kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	A375
细胞定位	Chromosome, Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human EHMT2/G9A.
序列	AAAAGAAAAAAEAGEAPAEMGALLEKETRGATERVHGSLGDTPRSEETLPKATPDSLEPAGPSSPASVTVTVGDEGADT PVGATPLIGDESENLEGDGDLRGGRILLGHATKSFPPSSPSKGGSCPSRAKMSMTGAGKSPPSVQSLAMRLLSMPGAQGA AAAGSEPPATTSPGQPKVHRARKTMSKPGNGQPPVPEKRPPEIQHFRMSDDVHSLGKVTSDLAKRRKLNSGGGLSEE LGSARRSGEVTLTKGDPGSLEEWEWTVVGDDFSLYDSYSVDERVSDSKSEVEALTEQLSE

靶点信息

研究背景	This gene encodes a methyltransferase that methylates lysine residues of histone H3. Methylation of H3 at lysine 9 by this protein results in recruitment of additional epigenetic regulators and repression of transcription. This gene was initially thought to be two different genes, NG36 and G9a, adjacent to each other in the HLA locus. Alternative splicing results in multiple transcript variants.
------	--

基因ID	10919
基因名	EHMT2
Swiss	Q96KQ7
别名	EHMT2; BAT8; C6orf30; G9A; GAT8; KMT1C; NG36; histone-lysine N-methyltransferase EHMT2

产品验证



Western blot analysis of EHMT2/G9A expressed in A375 using EHMT2/G9A Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn