

Argonaute 2 Rabbit mAb

货号: **AYM30281**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC IP FC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 IP: 1:20 - 1:50 FC: 1:20 - 1:50
理论分子量	93kDa/97kDa
实测分子量	97kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Jurkat,293T,HeLa
细胞定位	Cytoplasm,Nucleus,P-body
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human Argonaute 2.
序列	MDIPKIDIYHYELDIKPEKCPRRVNREIVEHMQHFKTQIFGDRKPVFDGRKNLYTAMPLPIGRDKVEL

靶点信息

研究背景	This gene encodes a member of the Argonaute family of proteins which play a role in RNA interference. The encoded protein is highly basic, and contains a PAZ domain and a PIWI domain. It may interact with dicer1 and play a role in short-interfering-RNA-mediated gene silencing. Multiple transcript variants encoding different isoforms have been found for this gene.
------	---

基因ID	27161
基因名	AGO2
Swiss	Q9UKV8
别名	AGO2;CASC7;EIF2C2;LINC00980;PPD;Q10;protein argonaute-2;Argonaute 2;EIF2C2

产品验证



Western blot analysis of Argonaute 2 expressed in Jurkat,293T,HeLa using Argonaute 2 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn