

Fukutin Rabbit mAb

货号: **AYM30286**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200
理论分子量	51kDa
实测分子量	51kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	BxPC-3,RD,U-87MG,Mouse testis,Mouse brain,Mouse heart,Rat liver
细胞定位	Golgi apparatus membrane,Single-pass type II membrane protein
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human Fukutin.
序列	DVKLDVFFFEETDHMWNGGTQAKTGKKFKYLFKFTLCWTEFVDMKVHVP CETLEYIEANYGKTWKIPVKTWDWKRSP PNVQPNGIWPISWDEVIQLY

靶点信息

研究背景	The protein encoded by this gene is a putative transmembrane protein that is localized to the cis-Golgi compartment, where it may be involved in the glycosylation of alpha-dystroglycan in skeletal muscle. The encoded protein is thought to be a glycosyltransferase and could play a role in brain development. Defects in this gene are a cause of Fukuyama-type congenital muscular dystrophy (FCMD), Walker-Warburg syndrome (WWS), limb-girdle muscular dystrophy type 2M (LGMD2M), and dilated cardiomyopathy type 1X (CMD1X). Alternatively spliced transcript variants have been found for this gene.
------	--

基因ID	2218
基因名	FKTN
Swiss	O75072
别名	FKTN; CMD1X; FCMD; LGMD2M; MDDGA4; MDDGB4; MDDGC4; fukutin

产品验证



Western blot analysis of Fukutin expressed in Mouse testis,Rat liver,BxPC-3 using Fukutin Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn