

HIP1 Rabbit mAb

货号: **AYM30295**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IF/ICC IP
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200 IP: 1:20 - 1:50
理论分子量	110kDa/113kDa/116kDa
实测分子量	116kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	A-549,BT-474,SW480,293T,Mouse brain
细胞定位	Cytoplasm,Cytoplasmic vesicle,Endomembrane system,Nucleus,clathrin-coated vesicle membrane
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human HIP1.
序列	SELEADLAEQQHLRQQAADDCEFLRAELDELRRQREDTEKAQRSLSEIERKAQANEQRYSKLKEKYSSELVQNHADLLRKN AEVTKQVSMARQAQVDLEREKKELEDSSLERISDQGQRKTQEQLVLES LKQELATSQRELQVLQGSLETSQAQSEANWAA EFAELEKERDSLVSAAHREEELSALRKELQDTQLKLASTEESMCQLAKDQRKMLLVGSRKAAEQVIQDALNQLPEEPLISG AGSADHLLSTVTSISSCIEQLEKSWSQYLACPEDISGLLH

靶点信息

研究背景	The product of this gene is a membrane-associated protein that functions in clathrin-mediated endocytosis and protein trafficking within the cell. The encoded protein binds to the huntingtin protein in the brain; this interaction is lost in Huntington's disease. Alternative splicing results in multiple transcript variants.
------	--

基因ID	3092
基因名	HIP1
Swiss	O00291
别名	HIP1;HIP-I;ILWEQ;SHON;SHONbeta;SHONgamma

产品验证



Western blot analysis of HIP1 expressed in Mouse brain,A-549,BT-474 using HIP1 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn