

— ABLYBIO, Help Your Research



# Phospho-ACC(S79) Rabbit mAb

货号: **AYM30297**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	
实测分子量	265kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	
细胞定位	Cytoplasm, cytosol
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein corresponding to mouse Phospho-ACC(S79) .
------	--

## 靶点信息

研究背景	
基因ID	3134

基因名	Acaca
Swiss	Q5SWU9 ( <a href="https://www.uniprot.org/uniprotkb/Q5SWU9/entry">https://www.uniprot.org/uniprotkb/Q5SWU9/entry</a> ), M ( <a href="https://www.uniprot.org/uniprotkb/M/entry">https://www.uniprot.org/uniprotkb/M/entry</a> ), Q13085H ( <a href="https://www.uniprot.org/uniprotkb/Q13085H/entry">https://www.uniprot.org/uniprotkb/Q13085H/entry</a> )
别名	Phospho-ACC(S79),Phospho-ACC(S79) Rabbit mAb,Acaca,ACC-alpha,Acetyl-CoA carboxylase 265,Acac,Gm 738

## 产品验证



Western blot analysis of Phospho-ACC(S79) expressed in human esophagus cancer, mouse lung, rat testis using Phospho-ACC(S79) Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn//www.ablybio.cn>)