

GBP1 Rabbit mAb

货号: **AYM30353**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	67kDa
实测分子量	68kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	5637(HTB-9)
细胞定位	Cell membrane,Cytoplasm,Cytoplasmic side,Golgi apparatus membrane,Lipid-anchor,Secreted
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human GBP1.
序列	ELAAQLEKKRDDFCQKQNEASSDRCSALLQVIFSPLEEEVKAGIYSKPGGYRLFVQKLQDLKKKYYEPRKGIQAEIILQTYL KSKESMTDAILQTDQTLTEKEKEIEVERVKAESAQASAKMLQEMQRKNEQMMEQKERSYQEHLKQLTEKMENDRVQLLK EQERTLALKLQEQEQLLKEGFQKESRIMKNEIQDLQTKMRRRKACTIS

靶点信息

研究背景	Guanylate binding protein expression is induced by interferon. Guanylate binding proteins are characterized by their ability to specifically bind guanine nucleotides (GMP, GDP, and GTP) and are distinguished from the GTP-binding proteins by the presence of 2 binding motifs rather than 3.
基因ID	2633

基因名	GBP1
Swiss	P32455
别名	GBP1

产品验证



Western blot analysis of GBP1 expressed in 5637(HTB-9) using GBP1 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn