

Phospho-SMAD5 (S463/S465) Rabbit mAb

货号: AYM30363

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200
理论分子量	52kDa
实测分子量	52kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	Rat lung
细胞定位	Cytoplasm, Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human Phospho-SMAD5 (S463/S465).
序列	STYPNSPASSGPGSPFQLPADTPPPAYMPPDDQMGQDNSQPMDSNNMIPQIMPSISSRDVQPVAAYEEPKHWCSIVYYEL NNRVGEAFHASSTSVLVDGFT

靶点信息

研究背景	The protein encoded by this gene is involved in the transforming growth factor beta signaling pathway that results in an inhibition of the proliferation of hematopoietic progenitor cells. The encoded protein is activated by bone morphogenetic proteins type 1 receptor kinase, and may be involved in cancer. Alternative splicing results in multiple transcript variants.
基因ID	4090

基因名	SMAD5
Swiss	Q99717
别名	DWFC;JV5-1;MADH5;SMAD5

产品验证



Western blot analysis of Phospho-SMAD5 (S463/S465) expressed in Rat lung using Phospho-SMAD5 (S463/S465) Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn