

Oct-1 Rabbit mAb

货号: **AYM30364**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200
理论分子量	95kDa
实测分子量	95kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,SGC-7901,Mouse testis
细胞定位	Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human Oct-1.
序列	MNNPSETSKPSMESGDGNTGTQTNGLDFQKQPVVGGAIATAQAQAFGLHGHVQLAGTSLQAAAQSLNVQSKSNEES GDSQQPSQPSQPSVQAAIPQTQLMLAGGQITGLTLTPAQQLLLQQAQAQQLLAAAVQQHSASQQHSAAGATISAS AATPMT

靶点信息

研究背景	The OCT1 transcription factor was among the first identified members of the POU transcription factor family (summarized by Sturm et al., 1993 [PubMed 8314572]). Members of this family contain the POU domain, a 160-amino acid region necessary for DNA binding to the octameric sequence ATGCAAAT.[supplied by OMIM, Jul 2010]
基因ID	5451

基因名	POU2F1
Swiss	P14859
别名	OCT1; OTF1; oct-1B

产品验证



Western blot analysis of Oct-1 expressed in HeLa,SGC-7901,Mouse testis using Oct-1 Rabbit mAb at 1:1 000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. B locking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn