

— ABLYBIO, Help Your Research



Ube2N / Ubc13 Rabbit mAb

货号: **AYM30369**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	17kDa
实测分子量	17kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HCT116,293T,C6,Mouse testis,Mouse brain,Rat testis,Rat brain
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human Ube2N / Ubc13.
------	------------------------------------------------------------------

靶点信息

研究背景	The modification of proteins with ubiquitin is an important cellular mechanism for targeting abnormal or short-lived proteins for degradation. Ubiquitination involves at least three classes of enzymes: ubiquitin-activating enzymes, or E1s, ubiquitin-conjugating enzymes, or E2s, and ubiquitin-protein ligases, or E3s. This gene encodes a member of the E2 ubiquitin-conjugating enzyme family. Studies in mouse suggest that this protein plays a role in DNA postreplication repair. [provided by RefSeq, Jul 2008]
基因ID	7334
基因名	UBE2N
Swiss	P61088
别名	Ube2N / Ubc13,Ube2N / Ubc13 Rabbit mAb,UBE2N,Bendless-like ubiquitin-conjugating enzyme,E2 ubiquitin-conjugating enzyme N,Ubc13,UbcH13,Ubiquitin carrier protein N,Ubiquitin-protein ligase N,BLU

产品验证



Western blot analysis of Ube2N / Ubc13 expressed in Mouse testis,Rat testis,HCT116 using Ube2N / Ubc13 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)