

— ABLYBIO, Help Your Research



HDAC8 Rabbit mAb

货号: **AYM30374**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC IP FC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 IP: 1:20 - 1:50 FC: 1:20 - 1:50
理论分子量	15kDa/16kDa/17kDa/28kDa/29kDa/31kDa/41kDa
实测分子量	42kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	C6
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human HDAC8 .
------	---

靶点信息

研究背景	Histones play a critical role in transcriptional regulation, cell cycle progression, and developmental events. Histone acetylation/deacetylation alters chromosome structure and affects transcription factor access to DNA. The protein encoded by this gene belongs to class I of the histone deacetylase family. It catalyzes the deacetylation of lysine residues in the histone N-terminal tails and represses transcription in large multiprotein complexes with transcriptional co-repressors. Multiple transcript variants encoding different isoforms have been found for this gene.
基因ID	55869
基因名	HDAC8
Swiss	Q9BY41
别名	HDAC8,HDAC8 Rabbit mAb,Protein deacetylase HDAC8,Protein deacetylase HDAC8,HDACL1

产品验证



Western blot analysis of HDAC8 expressed in C6 using HDAC8 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/>www.ablybio.cn)