

RBM3 Rabbit mAb

货号: **AYM30376**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC FC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 FC: 1:20 - 1:50
理论分子量	17kDa
实测分子量	17kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,Jurkat,HL-60
细胞定位	Cell projection,Cytoplasm,Nucleus,dendrite
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human RBM3.
序列	MSSEEGKLFVGGGLNFNTDEQALEDHFSSFGPISEVVVVKDRETQRSRGFGFITFTNPEHASVAMRAMNGESLDGRQIRV DHAGKSARGTRGGGFGAHGRGRSYSRGGDQGYGSGRYDYSRPGGYGYGYSRDYNGRNQGGYDRYSGGNYRD NYDN

靶点信息

研究背景	This gene is a member of the glycine-rich RNA-binding protein family and encodes a protein with one RNA recognition motif (RRM) domain. Expression of this gene is induced by cold shock and low oxygen tension . A pseudogene exists on chromosome 1. Multiple alternatively spliced transcript variants that are predicted to encode different isoforms have been characterized although some of these variants fit nonsense-mediated decay (NMD) criteria.
------	---

基因ID	5935
基因名	RBM3
Swiss	P98179
别名	RBM3;IS1-RNPL;RNPL

产品验证



Western blot analysis of RBM3 expressed in HeLa, Jurkat, HL-60 using RBM3 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn