

— ABLYBIO, Help Your Research



DPF2 Rabbit mAb

货号: **AYM30384**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC IP
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 IP: 1:20 - 1:50
理论分子量	23kDa/44kDa
实测分子量	44kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,Mouse thymus,Mouse testis,Rat testis,Rat lung
细胞定位	Nucleus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human DPF2.
------	---

靶点信息

研究背景	The protein encoded by this gene is a member of the d4 domain family, characterized by a zinc finger-like structural motif. This protein functions as a transcription factor which is necessary for the apoptotic response following deprivation of survival factors. It likely serves a regulatory role in rapid hematopoietic cell growth and turnover. This gene is considered a candidate gene for multiple endocrine neoplasia type I, a n inherited cancer syndrome involving multiple parathyroid, enteropancreatic, and pituitary tumors.
基因ID	5977
基因名	DPF2
Swiss	Q92785 (https://www.uniprot.org/uniprotkb/Q92785/entry)
别名	DPF2,DPF2 Rabbit mAb,Apoptosis response zinc finger protein,BRG1-associated factor 45D,D4,zinc and double PHD fingers family 2,Protein requiem,BAF45D,REQ,UBID4

产品验证



Western blot analysis of DPF2 expressed in Mouse thymus,Rat testis,HeLa using DPF2 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/>www.ablybio.cn)