

— ABLYBIO, Help Your Research



fetal hemoglobin Rabbit mAb

货号: **AYM30386**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	12kDa
实测分子量	12kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	K-562
细胞定位	cytosol,hemoglobin complex
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human fetal hemoglobin.
------	---

靶点信息

研究背景	The gamma globin genes (HBG1 and HBG2) are normally expressed in the fetal liver, spleen and bone marrow. Two gamma chains together with two alpha chains constitute fetal hemoglobin (HbF) which is normally replaced by adult hemoglobin (HbA) at birth. In some beta-thalassemias and related conditions, gamma chain production continues into adulthood. The two types of gamma chains differ at residue 136 where glycine is found in the G-gamma product (HBG2) and alanine is found in the A-gamma product (HBG1). The former is predominant at birth. The order of the genes in the beta-globin cluster is: 5'-epsilon -- gamma-G -- gamma-A -- delta -- beta--3'. [provided by RefSeq, Jul 2008]
基因ID	3047,3048
基因名	HBG1,HBG2
Swiss	P69891P69892 (https://www.uniprot.org/uniprotkb/P69891P69892/entry)
别名	fetal hemoglobin,fetal hemoglobin Rabbit mAb,HBG1,HBG2

产品验证



Western blot analysis of fetal hemoglobin expressed in K-562 using fetal hemoglobin Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)