— ABLYBIO, Help Your Research

Destrin Rabbit mAb

货号: AYM30391

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC IP FC
推荐浓度	**WB:** 1:500 - 1:2000 **IHC:** 1:50 - 1:200 **IF/ICC:** 1:50 - 1:200 **IP:** 1:20 - 1:50 **FC:** 1:20 - 1:50
理论分子量	17kDa
实测分子量	17kDa
形式	Liquid
保存条件	Store at -20℃. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	A-549,MCF7,NIH/3T3,C6,Mouse lung,Mouse spleen,Rat lung,Rat spleen
细胞定位	actin cytoskeleton,cortical actin cytoskeleton,cytoplasm,extracellular exosome
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human Destrin.

靶点信息

研究背景	The product of this gene belongs to the actin-binding proteins ADF family. This family of proteins is respo nsible for enhancing the turnover rate of actin in vivo. This gene encodes the actin depolymerizing protei n that severs actin filaments (F-actin) and binds to actin monomers (G-actin). Two transcript variants enc oding distinct isoforms have been identified for this gene. [provided by RefSeq, Jul 2008]
基因ID	11034
基因名	DSTN
Swiss	P60981 (https://www.uniprot.org/uniprotkb/P60981/entry)
别名	Destrin,Destrin Rabbit mAb,DSTN,Actin-depolymerizing factor,ACTDP,DSN

产品验证



Western blot analysis of Destrin expressed in Mouse lung,Rat lung,A-549 using Destrin Rabbit mAb at 1:1 000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blo cking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (https://www.ablybio.cn//www.ablybio.cn)