

Ctip2 Rabbit mAb

货号: **AYM30396**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	140kDa
实测分子量	140kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Jurkat,Raji,SH-SY5Y
细胞定位	neuron projection,nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human Ctip2.
序列	TGEKPYKCQLCDHACSQASKLKRHMKTHMHKAGSLAGRSDDGLSAASSPEPGTSELAGEGLKAADGDFRHHESDPSLG HEPEEEDEEEEEEEEELLENES

靶点信息

研究背景	This gene encodes a C2H2-type zinc finger protein and is closely related to BCL11A, a gene whose translocation may be associated with B-cell malignancies. Although the specific function of this gene has not been determined, the encoded protein is known to be a transcriptional repressor, and is regulated by the NURD nucleosome remodeling and histone deacetylase complex. Four alternatively spliced transcript variants encoding distinct isoforms have been found for this gene.
基因ID	64919

基因名	BCL11B
Swiss	Q9C0K0
别名	ATL1; ATL1-alpha; ATL1-beta; ATL1-delta; ATL1-gamma; CTIP-2; CTIP2; IMD49; RIT1; ZNF856B; hRIT1-alpha

产品验证



Western blot analysis of CtIP2 expressed in Jurkat, Raji, SH-SY5Y using CtIP2 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn