

TAF3 Rabbit mAb

货号: **AYM30404**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	103kDa
实测分子量	150kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	K-562,HeLa,mouse spleen,mouse brain,rat brain
细胞定位	male germ cell nucleus,nuclear membrane,nucleoplasm,nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human TAF3.
序列	HRYSELYGRTPILDDVGEAFQLMGVSLHELEDYIHNIIEPVTFFPHQIPSPVSKNNVLQFPQPGSKDAEERKEYIPDYLPPIVSSQEEEEEEQVPTDGGTS

靶点信息

研究背景	The highly conserved RNA polymerase II transcription factor TFIID (see TAF1; MIM 313650) comprises the TATA box-binding protein (TBP; MIM 600075) and a set of TBP-associated factors (TAFs), including TAF3. TAFs contribute to promoter recognition and selectivity and act as antiapoptotic factors.
基因ID	83860

基因名	TAF3
Swiss	Q5VWG9
别名	TAF140;TAFII140;TAFII-140;TAF3

产品验证



Western blot analysis of TAF3 expressed in K-562, HeLa, mouse spleen using TAF3 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn