

CHD3 Rabbit mAb

货号: **AYM30408**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	222KDa/226KDa/233KDa
实测分子量	260kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,Jurkat,Mouse brain,Rat brain
细胞定位	Cytoplasm,Nucleus,centrosome,cytoskeleton,microtubule organizing center
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human CHD3.
序列	MKAADTVILWARSKNDQLRISFPPGLCWGDRMPDKDDIRLLPSALGVKKRKRGPKKQKENKPGKPRKRKKRDSEEEFGS ERDEYREKSESGGSEYGTGPG

靶点信息

研究背景	This gene encodes a member of the CHD family of proteins which are characterized by the presence of chromatin organization modifier domains and SNF2-related helicase/ATPase domains. This protein is one of the components of a histone deacetylase complex referred to as the Mi-2/NuRD complex which participates in the remodeling of chromatin by deacetylating histones. Chromatin remodeling is essential for many processes including transcription. Autoantibodies against this protein are found in a subset of patients with dermatomyositis. Three alternatively spliced transcripts encoding different isoforms have been described. [provided by RefSeq, Jul 2008]
基因ID	1107
基因名	CHD3
Swiss	Q12873
别名	Mi-2a; Mi2-ALPHA; ZFH

产品验证



Western blot analysis of CHD3 expressed in Mouse brain, Rat brain, HeLa using CHD3 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn