— ABLYBIO, Help Your Research

Cytokeratin 10 Rabbit mAb

货号: AYM30424

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	**WB:** 1:500 - 1:2000 **IHC:** 1:50 - 1:200 **IF/ICC:** 1:50 - 1:200
理论分子量	60kDa
实测分子量	60kDa
形式	Liquid
保存条件	Store at -20℃. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,A-431
细胞定位	cell surface,cytoplasm,cytoskeleton,cytosol,extracellular exosome,extracellular space,nucleus
纯化	Affinity purification

抗原信息

抗原信息	Recombinant fusion protein corresponding to Human Cytokeratin 10.

靶点信息

研究背景	This gene encodes a member of the type I (acidic) cytokeratin family, which belongs to the superfamily of intermediate filament (IF) proteins. Keratins are heteropolymeric structural proteins which form the inter mediate filament. These filaments, along with actin microfilaments and microtubules, compose the cytos keleton of epithelial cells. Mutations in this gene are associated with epidermolytic hyperkeratosis. This g ene is located within a cluster of keratin family members on chromosome 17q21. [provided by RefSeq, Jul 2008]
基因ID	3858
基因名	KRT10
Swiss	P13645 (https://www.uniprot.org/uniprotkb/P13645/entry)
别名	Cytokeratin 10,Cytokeratin 10 Rabbit mAb,KRT10,Cytokeratin-10,Keratin-10,KPP

产品验证



Western blot analysis of Cytokeratin 10 expressed in HeLa,A-431 using Cytokeratin 10 Rabbit mAb at 1:10 00. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Block ing buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.



Immunohistochemical analysis of paraffin-embedded human bladder, using Cytokeratin 10 Antibody.

实验步骤

访问官网浏览详情: www.ablybio.cn (https://www.ablybio.cn//www.ablybio.cn)