

Cdk9 Rabbit mAb

货号: **AYM30441**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC IP
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 IP: 1:20 - 1:50
理论分子量	42kDa/53kDa
实测分子量	43kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,293T,MCF7,NIH/3T3,C6
细胞定位	Cytoplasm,Nucleus,PML body
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human Cdk9.
序列	ANVLITRDGVLKLDLADFLARAFSLAKNSQPNRYTNRVVTLWYRPELLLGERDYGPPIDLWGAGCIMAEMWTRSPIMQGN TEQHQALALISQLCGSITPEVWPVNDNYELYEKLELVKQGKRVKDRKAYVRDPYALDLIDKLLVLDPAQRIDSDDALNHDF FWSDPMPSDLKGMLSTHLTSMFEYLAPPRKGSQITQQSTNQSARNPATTNQTQEFERVF


靶点信息

研究背景	The protein encoded by this gene is a member of the cyclin-dependent protein kinase (CDK) family. CDK family members are highly similar to the gene products of <i>S. cerevisiae</i> cdc28, and <i>S. pombe</i> cdc2, and known as important cell cycle regulators. This kinase was found to be a component of the multiprotein complex TAK/P-TEFb, which is an elongation factor for RNA polymerase II-directed transcription and functions by phosphorylating the C-terminal domain of the largest subunit of RNA polymerase II. This protein forms a complex with and is regulated by its regulatory subunit cyclin T or cyclin K. HIV-1 Tat protein was found to interact with this protein and cyclin T, which suggested a possible involvement of this protein in AIDS.
基因ID	1025
基因名	CDK9
Swiss	P50750
别名	CDK9;C-2k;CDC2L4;CTK1;PITALRE;TAK

产品验证



Immunohistochemical analysis of paraffin-embedded human cervix cancer, using CDK9 Antibody.



Western blot analysis of Cdk9 expressed in HeLa, 293T, MCF7 using Cdk9 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 1 20s.

实验步骤

访问官网浏览详情: www.ablybio.cn