

CCDC98 Rabbit mAb

货号: **AYM30457**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IF/ICC IP FC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200 IP: 1:20 - 1:50 FC: 1:20 - 1:50
理论分子量	50kDa
实测分子量	50kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,MCF7,SKOV3
细胞定位	Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human CCDC98.
序列	SCNYNHHLDVVDNLTLMVEHTDIPEASPASTPQIIKHKALDLDLDRWQFKRSRLDLDQDKRSKADTGSSNQDKASKMSSPETDEEIEKMKGFGEYSRSPTF

靶点信息

研究背景	This gene encodes a protein that binds to the C-terminal repeats of breast cancer 1 (BRCA1) through a phospho-SXXF motif. The encoded protein recruits ubiquitin interaction motif containing 1 protein to BRCA1 protein and is required for DNA damage resistance, DNA repair, and cell cycle checkpoint control. Pseudo genes of this gene are found on chromosomes 3 and 8. Alternative splicing results in multiple transcript variants.
------	--

基因ID	84142
基因名	ABRAXAS1
Swiss	Q6UWZ7
别名	FAM175A; ABRA1; CCDC98; BRCA1-A complex subunit Abraxas

产品验证



Western blot analysis of CCDC98 expressed in 293T, MCF7, SKOV3 using CCDC98 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn