

# AZI2 Rabbit mAb

货号: **AYM30470**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IF/ICC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IF/ICC:</b> 1:50 - 1:200
理论分子量	26kDa/28kDa/44kDa
实测分子量	45kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HepG2,SH-SY5Y,Mouse testis,Rat testis
细胞定位	cytoplasm
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein corresponding to Human AZI2.
序列	

## 靶点信息

研究背景	AZI2, or NAP1, contributes to the activation of NF $\kappa$ B (see MIM 164011)-dependent gene expression by activating IKK-related kinases, such as NAK (TBK1; MIM 604834) (Fujita et al., 2003 [PubMed 14560022]).[supplied by OMIM, Mar 2008]
基因ID	64343
基因名	AZI2

<b>Swiss</b>	Q9H6S1
别名	AZ2;NAP1;TILP;AZI2

## 产品验证



Western blot analysis of AZI2 expressed in Mouse testis,Rat testis,HepG2 using AZI2 Rabbit mAb at 1:10 00. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blo cking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)