

NUR77 Rabbit mAb

货号: **AYM30490**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	64kDa
实测分子量	64kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HepG2,SKOV3,U-87MG
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human NUR77.
序列	MPCIQAQYGTAPSPGPRDHLASDPLTPEFIKPTMDLASPEAAPAAPALPSFSTFMDGYTGEFDTFLYQLPGTVQPCSSA SSSASSTSSSSATSPASAS

靶点信息

研究背景	This gene encodes a member of the steroid-thyroid hormone-retinoid receptor superfamily. Expression is induced by phytohemagglutinin in human lymphocytes and by serum stimulation of arrested fibroblasts. The encoded protein acts as a nuclear transcription factor. Translocation of the protein from the nucleus to mitochondria induces apoptosis. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jan 2011]
基因ID	3164

基因名	NR4A1
Swiss	P22736
别名	GFRP1; HMR; N10; NAK-1; NGFIB; NP10; NUR77; TR3

产品验证



Western blot analysis of NUR77 expressed in HepG2,SKOV3,U-87MG using NUR77 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn