

CD5 Rabbit mAb

货号: **AYM30492**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC ICC IP
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 ICC: 1:50 - 1:200 IP: 1:20 - 1:50
理论分子量	54kDa
实测分子量	55kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Raji,Mouse heart,Mouse thymus
细胞定位	Cell membrane,Single-pass type I membrane protein
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human CD5.
序列	RLQLVAQSGGQHCAGVVEFYSGSLGGTISYEAQDKTQDLENFLC>NNLQCGSFLKHLPETEAGRAQDPGEPREHQPLPIQ WKIQNSSCTSLEHCFRKIKPKQKSGRVLALLCSGFQPKVQSRLVGGSSICEGTVEVRQGAQWAALCDSSSARSSLRWEEV CREQQCGSVNSYRVLDAGDPTSRGLFCPHQKLSQCHELWERNYSYCKKVFVTCQDPN

靶点信息

研究背景	This gene encodes a member of the scavenger receptor cysteine-rich (SRCR) superfamily. Members of this family are secreted or membrane-anchored proteins mainly found in cells associated with the immune system. This protein is a type-I transmembrane glycoprotein found on the surface of thymocytes, T lymphocytes and a subset of B lymphocytes. The encoded protein contains three SRCR domains and may act as a receptor to regulate T-cell proliferation. Alternative splicing results in multiple transcript variants encoding different isoforms.
基因ID	921
基因名	CD5
Swiss	P06127
别名	CD5;LEU1;T1

产品验证



Western blot analysis of CD5 expressed in Raji, Mouse heart, Mouse thymus using CD5 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn