

Tenascin C Rabbit mAb

货号: **AYM30553**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	
实测分子量	241kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-138MG
细胞定位	basement membrane,endoplasmic reticulum lumen,extracellular region,extracellular space,focal adhesion
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human Tenascin C.
序列	NGDTTSGLYTIYLNQDKAEALEVFCDMTSDGGGWIVFLRRKNGRENFYQNWKAYAAGFGDRREEFWLGLDNLNKITAQ GQYELRVDLRDHGETAFVYDKFSVGDATRYKLVKVEGYSGTAGDSMAYHNGRSFSTFDKDTDSAITNCALS YKGAFW YRNCHRVNLMGRYGDNNHSQGVNWFHWKGHEHSIQFAEMKLRPSNFRNLEGRKRRA

靶点信息

研究背景	This gene encodes an extracellular matrix protein with a spatially and temporally restricted tissue distribution. This protein is homo-hexameric with disulfide-linked subunits, and contains multiple EGF-like and fibronectin type-III domains. It is implicated in guidance of migrating neurons as well as axons during development, synaptic plasticity, and neuronal regeneration. [provided by RefSeq, Jul 2011]
基因ID	3371

基因名	TNC
Swiss	P24821
别名	150-225;DFNA56;GMEM;GP;HXB;JI;TN;TN-C;TNC

产品验证



Western blot analysis of Tenascin C expressed in U-138MG using Tenascin C Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn