

CLIP170 Rabbit mAb

货号: **AYM30585**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200
理论分子量	156kDa/160kDa/162kDa
实测分子量	130,170kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-87MG,Mouse heart
细胞定位	Cell projection,Cytoplasm,Cytoplasmic side,Cytoplasmic vesicle membrane,Peripheral membrane protein, cytoskeleton,ruffle
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human CLIP170.
序列	SALLTEKDAELEKLRNEVTVLRGENASAKSLHSVVTLES DKVKLELKVKNLELQLKENKRQLSSSSGNTDTQAEDEDERAQ ESQIDFLNSVIVDLQRKNQDLKMKVEMMSEAALNGNGDDLNNYDSDQEQSKKKPRLFCDICDFDLHDTEDCPTQA QMSEDPHSTHHGSRGEERPYCEICEMFGHWATNCNDDETF

靶点信息

研究背景	The protein encoded by this gene links endocytic vesicles to microtubules. This gene is highly expressed in Reed-Sternberg cells of Hodgkin disease. Several transcript variants encoding different isoforms have been found for this gene.
基因ID	6249

基因名	CLIP1
Swiss	P30622
别名	CLIP1;CLIP;CLIP-170;CLIP170;CYLN1;RSN

产品验证



Western blot analysis of CLIP170 expressed in U-87MG, Mouse heart using CLIP170 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn