

PGDH Rabbit mAb

货号: **AYM30590**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC FC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 FC: 1:20 - 1:50
理论分子量	15kDa/19kDa/21kDa/28kDa
实测分子量	29kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	A-549,THP-1(Negative control)
细胞定位	Cytoplasm
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human PGDH.
序列	MHVNGKVALVTGAAQIGRAFAEALLLKGAQVALVDWNLEAGVQCKAALDEQFEPQKTLFIQCDVADQQQLRDTFRKVV DHFGRLDILVNNAGVNNNEKNWEKTLQINLVSVISGTYLGLDYMSKQNGGEGGIIINMSSLAGLMPVAQQPVYCASKHGIV GFTRSAALANLMNSGVRLNAICPGFVNTAILESIEKEENMGQYIEYKDHDKMIKYYGILDPPLIANGLITLIEDDALNGAIMKI TTSKGIHFQDYDTTPFQAKTQ

靶点信息

研究背景	This gene encodes a member of the short-chain nonmetalloenzyme alcohol dehydrogenase protein family . The encoded enzyme is responsible for the metabolism of prostaglandins, which function in a variety of physiologic and cellular processes such as inflammation. Mutations in this gene result in primary autosomal recessive hypertrophic osteoarthropathy and craniosteoarthropathy. Multiple transcript variants encoding different isoforms have been found for this gene.
基因ID	3248
基因名	HPGD
Swiss	P15428
别名	HPGD;15-PGDH;PGDH;PGDH1;PHOAR1;SDR36C1

产品验证



Western blot analysis of PGDH expressed in A-549,THP-1(Negative control) using PGDH Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn