

ANGPTL3 Rabbit mAb

货号: **AYM30596**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200
理论分子量	54kDa
实测分子量	54kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HepG2,A375,Mouse lung,Mouse liver,Rat lung,Rat liver
细胞定位	Secreted
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human ANGPTL3.
序列	MFTIKLLLIVPLVISSRIDQDNSSFDLSPEPKSRFAMLDDVKILANGLLQLGHGLKDFVHKTGQINDIFQKLNIFDQSFYD LSLQTSEIKEEEKELR

靶点信息

研究背景	This gene encodes a member of a family of secreted proteins that function in angiogenesis. The encoded protein, which is expressed predominantly in the liver, is further processed into an N-terminal coiled-coil domain-containing chain and a C-terminal fibrinogen chain. The N-terminal chain is important for lipid metabolism, while the C-terminal chain may be involved in angiogenesis. Mutations in this gene cause familial hypobetalipoproteinemia type 2. [provided by RefSeq, Aug 2015]
基因ID	27329

基因名	ANGPTL3
Swiss	Q9Y5C1
别名	ANG-5; ANGPT5; ANL3; FHBL2

产品验证



Western blot analysis of ANGPTL3 expressed in Mouse lung,Rat lung,HepG2 using ANGPTL3 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120 S.

实验步骤

访问官网浏览详情: www.ablybio.cn