

TRIM25 Rabbit mAb

货号: **AYM30606**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC IP FC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 IP: 1:20 - 1:50 FC: 1:20 - 1:50
理论分子量	70kDa
实测分子量	71kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,293T,MCF7
细胞定位	Cytoplasm
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human TRIM25.
序列	SPNAQVACDHCLKEAAVKTCLVCMASFCQEHLQPHFDSPAQDHPQLPPVRDLLRRKCSQHNRLREFFCPEHSECICHICLVEHKTCSPASLSQASADLEATLRHKLTVMYSQINGASRALDDVRNRQQDVRMTANRKVEQLQQEYTEMKALLDASETTS TRKIKKEEKRVNSKFDTIYQILLKKKSEIQTLKEEIEQSLTKRDEFEFLEKASKLRGISTKPVYIPEVELNHKLIKGIHQSTIDLKN ELKQCIGRLQEPTPSSGDPGEHDPASTHKSTRPVKKVSKEEKSKKPPPPALP

靶点信息

研究背景	The protein encoded by this gene is a member of the tripartite motif (TRIM) family. The TRIM motif includes three zinc-binding domains, a RING, a B-box type 1 and a B-box type 2, and a coiled-coil region. The protein localizes to the cytoplasm. The presence of potential DNA-binding and dimerization-transactivation domains suggests that this protein may act as a transcription factor, similar to several other members of the TRIM family. Expression of the gene is upregulated in response to estrogen, and it is thought to mediate estrogen actions in breast cancer as a primary response gene.
基因ID	7706
基因名	TRIM25
Swiss	Q14258
别名	TRIM25;EFP;RNF147;Z147;ZNF147

产品验证



Western blot analysis of TRIM25 expressed in HeLa,293T,MCF7 using TRIM25 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn