

AAMP Rabbit mAb

货号: **AYM30613**

产品信息

| | |
|-------|---|
| 反应 | Human,Mouse,Rat |
| 宿主 | Rabbit |
| 克隆性 | Monoclonal |
| 预测反应 | |
| 应用 | WB IF/ICC FC |
| 推荐浓度 | WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200 FC: 1:20 - 1:50 |
| 理论分子量 | 46kDa |
| 实测分子量 | 52kDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | 293T,MCF7,Mouse liver,Mouse testis,Rat liver,Rat testis,A375 |
| 细胞定位 | Cell membrane,Cytoplasm |
| 纯化 | Affinity purification |

抗原信息


| | |
|------|--|
| 抗原信息 | Recombinant fusion protein corresponding to Human AAMP. |
| 序列 | MESESESGAAADTPPLETLSFHGDEEIIIEVVVLDPGPPDPPDLAQEMEDVDFEEEEEEGNEEGWVLEPQEGVVGSMEGP DDSEVTFALHSASVFCVSLD |

靶点信息

| | |
|------|--|
| 研究背景 | The gene is a member of the immunoglobulin superfamily. The encoded protein is associated with angiogenesis, with potential roles in endothelial tube formation and the migration of endothelial cells. It may also regulate smooth muscle cell migration via the RhoA pathway. The encoded protein can bind to heparin and may mediate heparin-sensitive cell adhesion. |
| 基因ID | 14 |

| | |
|-------|--------|
| 基因名 | AAMP |
| Swiss | Q13685 |
| 别名 | AAMP |

产品验证



Western blot analysis of AAMP expressed in Mouse liver,Rat liver,293T using AAMP Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn