

UAP56 Rabbit mAb

货号: **AYM30614**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC FC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 FC: 1:20 - 1:50
理论分子量	48kDa/50kDa
实测分子量	45kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,HeLa,LO2,SKOV3,K-562,SW620
细胞定位	Cytoplasm,Nucleus,Nucleus speckle
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human UAP56.
序列	RGMPQEERLSRYQQFKDFQRRILVATNLFGRGMDIERNVIAFNVDMPEDSDTYLHRVARAGRFGTKGLAITFVSDENDAKI LNDVQDRFEVNISELPDEIDISSYIEQTR

靶点信息

研究背景	This gene encodes a member of the DEAD box family of RNA-dependent ATPases that mediate ATP hydrolysis during pre-mRNA splicing. The encoded protein is an essential splicing factor required for association of U2 small nuclear ribonucleoprotein with pre-mRNA, and it also plays an important role in mRNA export from the nucleus to the cytoplasm. This gene belongs to a cluster of genes localized in the vicinity of the genes encoding tumor necrosis factor alpha and tumor necrosis factor beta. These genes are all within the human major histocompatibility complex class III region. Mutations in this gene may be associated with rheumatoid arthritis. Alternative splicing results in multiple transcript variants. Related pseudogenes have been identified on both chromosomes 6 and 11. Read-through transcription also occurs between this gene and the upstream ATP6V1G2 (ATPase, H ⁺ transporting, lysosomal 13kDa, V1 subunit G2) gene.
基因ID	7919
基因名	DDX39B
Swiss	Q13838
别名	DDX39B;BAT1;D6S81E;UAP56

产品验证



Western blot analysis of UAP56 expressed in 293T, HeLa, LO2 using UAP56 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn