

— ABLYBIO, Help Your Research



Cdk7 Rabbit mAb

货号: **AYM30629**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	39kDa
实测分子量	39kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T
细胞定位	Cytoplasm,Nucleus,perinuclear region
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human Cdk7.
------	---

靶点信息

研究背景	The protein encoded by this gene is a member of the cyclin-dependent protein kinase (CDK) family. CDK family members are highly similar to the gene products of <i>Saccharomyces cerevisiae</i> cdc28, and <i>Schizosaccharomyces pombe</i> cdc2, and are known to be important regulators of cell cycle progression. This protein forms a trimeric complex with cyclin H and MAT1, which functions as a Cdk-activating kinase (CAK). It is an essential component of the transcription factor TFIIF, that is involved in transcription initiation and DNA repair. This protein is thought to serve as a direct link between the regulation of transcription and the cell cycle.
基因ID	1022
基因名	CDK7
Swiss	P50613
别名	Cdk7,Cdk7 Rabbit mAb,39 kDa protein kinase,CDK-activating kinase 1,Cell division protein kinase 7,Serine/threonine-protein kinase 1,TFIIF basal transcription factor complex kinase subunit,CAK,CAK1,CDKN7,M O15,STK1

产品验证



Western blot analysis of Cdk7 expressed in 293T using Cdk7 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/>www.ablybio.cn)